

Does Childhood Diarrhea Influence Cognition Beyond the Diarrhea-Stunting Pathway?

Christa L. Fischer Walker^{1*}, Laura Lamberti¹, Linda Adair², Richard L. Guerrant³, Andres G. Lescano^{4,5}, Reynaldo Martorell⁶, Relana C. Pinkerton³, Robert E. Black¹

1 Department of International Health, Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland, United States of America, **2** Department of Nutrition, University of North Carolina, Chapel Hill, North Carolina, United States of America, **3** Center for Global Health, University of Virginia, Charlottesville, Virginia, United States of America, **4** Department of Parasitology, U.S. Naval Medical Research Unit No. 6 (NAMRU-6), Lima, Peru, **5** School of Public Health and Management, Universidad Peruana Cayetano Heredia, Lima, Peru, **6** Hubert Department of Global Health, Rollins School of Public Health, Emory University, Atlanta, Georgia, United States of America

Abstract

Background: Diarrhea is a leading cause of morbidity among children under 5 years of age in low- and middle-income countries yet the additional effects and sequelae, such as cognitive impairment associated with diarrhea, have not been quantified.

Methods: We quantified the association between diarrhea prevalence and cognitive outcomes while controlling for linear growth in 4 study populations. Cognition was assessed using different methods across sites and was expressed in standardized units. We built linear regression models for each study with standardized cognitive score as the outcome and diarrhea prevalence as the main predictor variable. We then conducted meta-analyses of the regression coefficients to generate pooled estimates of the association between diarrhea prevalence and cognition whilst controlling for anthropometric status and other covariates.

Results: Diarrhea was not a significant predictor of cognitive score in any site in the regression models or in the meta-analyses (Coefficient = 0.07; 95% CI: -0.1, 0.2). The length for age Z- score was negatively related to cognition in all sites (0.18; 95% CI: 0.14, 0.21), with coefficients remarkably similar across sites (Coefficient Range: 0.168–0.186).

Conclusions: We did not demonstrate an association between diarrhea and cognition with stunting included in the model. The links between diarrhea, stunting, and cognition provide additional rationale for accelerating interventions to reduce diarrhea.

Citation: Fischer Walker CL, Lamberti L, Adair L, Guerrant RL, Lescano AG, et al. (2012) Does Childhood Diarrhea Influence Cognition Beyond the Diarrhea-Stunting Pathway? PLoS ONE 7(10): e47908. doi:10.1371/journal.pone.0047908

Editor: Thomas Eisele, Tulane University School of Public Health and Tropical Medicine, United States of America

Received: May 31, 2012; **Accepted:** September 17, 2012; **Published:** October 31, 2012

This is an open-access article, free of all copyright, and may be freely reproduced, distributed, transmitted, modified, built upon, or otherwise used by anyone for any lawful purpose. The work is made available under the Creative Commons CC0 public domain dedication.

Funding: This work was supported by multiple donors. Dr. Fischer Walker, Laura Lamberti, and D. Robert Black were funded by the Bill and Melinda Gates Foundation via the U.S. Fund for UNICEF for the Child Health Epidemiology Reference Group (CHERG). The participation of Dr. Lescano in this project is sponsored by the training grant NIH/FIC 2D43 TW007393 awarded to NAMRU-6 by the Fogarty International Center of the U.S. National Institutes of Health. The Brazil work was supported by NIH/NIAID ICIDR Grant No. U01AI026512. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: cfischer@jhsph.edu

Introduction

Diarrhea is a leading cause of morbidity and mortality among children under 5 years of age in low- and middle-income countries [1]. While the most severe diarrhea sequela, child mortality, is easily quantifiable, the indirect effects of diarrhea on subsequent risk of infection, growth, and cognitive ability are often less obvious and thus the full disease burden attributable to diarrhea is not as well understood [2].

Frequent diarrhea during the first 2 years of life has been negatively correlated with cognitive development and early school performance [3,4], yet the pathway for this association is not fully understood. In an analysis of 9 data sets, Checkley et al. estimated that each episode of diarrhea in early childhood increases the odds of being stunted at 24 mo of age by 5 [5]. In addition, it has also been shown that early childhood stunting contributes to cognitive

impairment [6,7], thus it is possible that diarrhea may be associated with a portion of cognitive impairment via the well-documented nutrition and infection cycle.

There is also evidence suggesting that cognitive delays may be related to differences in caregiver behavior toward a sickly child. During the acute and convalescent period of an illness, the child may be listless and less eager to play or explore his/her environment [8]. Children with prolonged or frequent episodes of diarrhea spend an increased proportion of days in the acute or recovery phase of illness and thus are likely to receive less stimulation by their own activity or by caregivers during these periods [8,9].

We sought to quantify the association of diarrhea on cognitive outcomes while controlling for stunting in 4 datasets from 4 distinct study populations. Published studies have found an association between diarrhea and cognition, but often do not take

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 31 OCT 2012		2. REPORT TYPE		3. DATES COVERED 00-00-2012 to 00-00-2012	
4. TITLE AND SUBTITLE Does Childhood Diarrhea Influence Cognition Beyond the Diarrhea-Stunting Pathway?				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Johns Hopkins Bloomberg School of Public Health, Department of International Health, 615 North Wolfe Street, Baltimore, MD, 21205				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT Background: Diarrhea is a leading cause of morbidity among children under 5 years of age in low- and middle-income countries yet the additional effects and sequelae, such as cognitive impairment associated with diarrhea, have not been quantified. Methods: We quantified the association between diarrhea prevalence and cognitive outcomes while controlling for linear growth in 4 study populations. Cognition was assessed using different methods across sites and was expressed in standardized units. We built linear regression models for each study with standardized cognitive score as the outcome and diarrhea prevalence as the main predictor variable. We then conducted meta-analyses of the regression coefficients to generate pooled estimates of the association between diarrhea prevalence and cognition whilst controlling for anthropometric status and other covariates. Results: Diarrhea was not a significant predictor of cognitive score in any site in the regression models or in the metaanalyses (Coefficient = 0.07; 95% CI: 20.1, 0.2). The length for age Z- score was negatively related to cognition in all sites (0.18; 95% CI: 0.14, 0.21), with coefficients remarkably similar across sites (Coefficient Range: 0.168?0.186). Conclusions: We did not demonstrate an association between diarrhea and cognition with stunting included in the model. The links between diarrhea, stunting, and cognition provide additional rationale for accelerating interventions to reduce diarrhea.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 6	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

into consideration anthropometric status [3,4]. By including only datasets with individual level data on diarrhea prevalence, anthropometry at 2 years of age, and subsequent cognitive outcomes we are able to consider the relationship of diarrhea to cognition distinct from associations observed via the stunting pathway.

Methods

Selection of Data Sets

We conducted a PubMed search to identify possible studies for inclusion using the following search terms: *diarrhea*, *stunting*, *length-for-age*, *growth*, *cognitive impairment*, *cognition*, and *children*. We searched for studies that conducted routine diarrhea surveillance among populations of children under 2 years of age for at least 1 year, collected anthropometric indicators and conducted a follow-up of cognitive development at least 1 year after the initial diarrhea surveillance period. We excluded studies that did not use a published methodology for assessing cognition.

Regression analysis

We sought to determine if diarrhea prevalence between 0 and 2 years of age had an impact on cognitive development later in life controlling for stunting at 2 years. To do this we built linear regression models for each study and subsequently conducted meta-analyses to pool each of the beta coefficients produced by these regressions. In the selected studies, cognitive scores were reported using 4 different methodologies and scales: non-verbal IQ [7,10], Test of Non-verbal Intelligence III (TONI III) [11], Wechsler Intelligence Scale for Children-Revised [12], and the first factor of a factor analysis of eight tests representing general perceptual-organizational and verbal abilities [13] in Philippines, Brazil, Peru, and Guatemala, respectively (Table 1). To account for the use of different cognitive scales across study sites, we generated standardized scores by subtracting the mean and dividing by the standard deviation.

For each study, we fitted two regression models with the standardized cognitive score as the outcome and diarrhea prevalence, defined as the percentage of days/study periods in which the caregiver reported diarrhea, as the main predictor variable. Model 1 controlled for length-for-age (LAZ) at age 2 as a continuous variable and model 2 controlled for LAZ as a categorical variable (with LAZ ≤ -2 as a the cut off for stunted vs. not stunted). LAZ was defined by the World Health Organization reference standard [14]. In both models, we also controlled for variables commonly thought to influence cognitive function and those found to be significant in bivariate models: sex, mother's education, age at cognitive evaluation, and socioeconomic status (SES). With the exception of sex and LAZ, the remaining covariates were not treated uniformly across regression analyses. Variation in variable definitions could not be avoided because all studies were uniquely designed and not for the purpose of combining data. SES was defined as log household income and asset ownership in the Philippines; monthly income and the number of household rooms per person in Brazil; log household income in Peru; and multi-component household SES score (through factor analysis) and village residence in Guatemala. Age at cognitive evaluation was recorded in months and defined as a continuous variable in the Philippines, Brazil and Peru; in Guatemala, children were four years of age at cognitive assessment and thus categorical indicator variables for birth year were included in the model. In the Philippines, Peru and Guatemala, maternal education was defined as the average number of years enrolled in school; whereas, in Brazil, maternal education was a categorical variable representing the percentage of mothers that had not completed primary school. While the covariates included in each of the four regression models were not identical, all models controlled for each of these defined variables.

Meta-analysis. We conducted meta-analyses to pool regression beta coefficients for variables with exact definitions across all four regression models (i.e. diarrhea prevalence, LAZ, and sex). In order to generate pooled estimates of regression beta coefficients, we used Comprehensive Meta Analysis (CMA) software, which

Table 1. Description of key study variables and variation in cognitive testing strategies across studies.

	Guatemala	Philippines	Brazil	Peru
Sample size	252	2117	111	143
Sex (% boys)	54.8	52.9	40.5	53
Stunted (% < -2 LAZ)	86.5	67.5	24.3	32
Maternal education (mean years enrolled in school)^a or % of mothers who had not completed primary school^b	1.4 ^a	7.0 ^a	81.1% ^b	7.3 ^a
Cognitive Follow-up				
Age at cognitive test, mean (range) in years	4 (± 1 week)	8.5 (8.4–8.7)	8 (6–12)	9.4 (8.4–10.1)
Type of cognitive test	First factor score of eight tests	Non-verbal IQ	TONI IQ	Wechsler (WISC-R)
Overview of cognitive test	Perceptual-organizational and verbal abilities	Cognitive test designed to assess fluid ability (i.e. analytic or reasoning skills)	Non-verbal intelligence, aptitude, abstract reasoning and problem solving. Language free validation in non-English speaking groups.	Three separate IQs calculated from 10 subtests, a Verbal Scale IQ, a Performance Scale IQ, and a Full Scale IQ

^aMaternal education expressed as mean years enrolled in school.

^bMaternal education expressed as % of mothers who had not completed primary school.

doi:10.1371/journal.pone.0047908.t001

requires the coefficient and its standard error [15]. We used STATA 11 to determine the z-statistic corresponding to the p-value of each coefficient and then estimated the standard error by dividing the coefficient by its z-statistic [16]. We conducted random effects meta-analyses for each beta coefficient to account for both the within and between study variation that is expected with data from four different study sites. We also plotted a funnel plot to look for possible bias among the included studies.

Results

We identified four studies that met our inclusion and exclusion criteria. The studies were conducted among 4 pediatric populations in low-income settings with ongoing diarrhea surveillance; details of the studies have been previously published [4,13,17,18,19,20,21]. In brief, children in these populations have high rates of infectious disease and are of low socioeconomic status. Diarrhea rates ranged from 4% of days in Brazil to 19% of days in The Philippines [4,13,17,18,19,20,21]. Table 1 describes the sample size and key study variables included in the analysis.

We conducted linear regression models for each study site to determine the within study effect, if any, of diarrhea on cognition. Diarrhea was not a significant predictor of cognitive score in any site (Table 2). The LAZ score coefficient was consistent across the 4 study populations and associated with cognition such that lower LAZ scores were related to lower cognitive performance (Coefficient Range: 0.168–0.186) in these multivariate analyses.

We then conducted meta-analyses to pool the regression coefficients across study sites and thus determine the cross-country effect of diarrhea on cognition whilst controlling for anthropometric status. The results of the two models were similar; here we present the pooled results of model 1 which considered length-for-age Z score as a continuous variable (Table 3). In pooled analyses, there was no statistically significant association between diarrhea and cognitive development (Coefficient = 0.07; 95% CI: −0.1, 0.2) (Figure 1). The symmetry observed in the funnel plot figure suggests that none of the included studies are biasing the results (Figure 2).

Discussion

We conducted a meta-analysis to assess the relationship between diarrhea and cognition while controlling for LAZ. We included 4 data sets from 4 different study populations and did not see an overall effect of diarrhea prevalence rates during the first 2 years of life on cognitive scores several years later. We did not observe an association of diarrhea prevalence and cognitive scores yet we did observe a remarkably consistent association of LAZ score and cognitive development across data sites. Thus, in these data sets a portion of cognitive impairment might be explained by the child's length deficit and this is independent of the child's diarrhea prevalence rate. It may be possible that diarrhea is a contributing factor to childhood stunting in these low-income settings but in this setting diarrhea prevalence alone did not appear to be associated with poorer cognitive performance in these children.

Some studies have observed associations between increasing diarrhea prevalence rates during early childhood and deficits or delays in later cognition [3,4]. However, these studies have not taken into account the child's anthropometric status at 2 years. Checkley and colleagues found diarrhea prevalence to account for 20% of the stunting prevalence, and thus one might suspect that diarrhea may account for some portion of the stunting-cognition relationship [5].

Our analysis has several potential methodological weaknesses. The studies included here assessed cognition with 4 distinct methodologies. These studies were not designed with this meta-analysis in mind; rather, they were designed to use the best-standardized cognitive test available, which could be locally adapted to each setting. We accounted for the differences in scoring practices by using a statistically standardized score for each regression; this enabled us to include all study sites in the meta-analysis, even though the outcome variables were scaled differently. The studies included here also had variable follow-up time and did not have standard socio-economic covariates. Though we accounted for these differences analytically, the analysis would have been stronger if these covariates were standard across the datasets.

We included age in the regression model for all countries. Although cognitive test scores are age standardized, the standardization method differs among tests, and thus including this variable

Table 2. Results of country-level linear regression models to determine the association of diarrhea on cognition.^{a,b,c,d,e}

	Philippines			Brazil			Peru			Guatemala		
	Coefficient	SE	p value	Coefficient	SE	p value	Coefficient	SE	p value	Coefficient	SE	p value
% days/periods with diarrhea	0.158	0.130	0.225	−2.900	2.162	0.183	1.207	1.277	0.347	0.030	0.070	0.700
LAZ (Continuous Variable)	0.177	0.019	0.000	0.186	0.081	0.024	0.168	0.092	0.070	0.170	0.060	0.000
Sex (0 = female, 1 = male)	−0.084	0.040	0.034	−0.155	0.187	0.409	−0.318	0.195	0.106	−0.210	0.130	0.110

^a. For each data set, standardized cognitive Z-score served as the outcome of linear regression.

^b. All four regression analyses controlled for diarrhea prevalence, stunting and sex in the same way. Diarrhea prevalence was coded as the percentage of days/periods during which diarrhea was reported; LAZ was treated as a continuous z-score; sex was a categorical variable (0 = male, 1 = female).

^c. SES was controlled for differently in each country given the data available from each site: 1) Philippines: controlled for log household income and ownership of assets; 2) Brazil: controlled for monthly income and the number of rooms in the household per person; 3) Peru: controlled for log household income; and 4) Guatemala: controlled for a multi-component SES score (through factor analysis) for the household as a continuous variable and residence in one of four villages.

^d. Age (in months) at cognitive evaluation was treated as a continuous variable in Philippines, Brazil, and Peru. In Guatemala, the children were 4 years of age at cognitive assessment, and therefore age was controlled for by the addition of categorical indicator variables for birth year to the regression model.

^e. In Philippines, Peru and Guatemala, maternal education was defined as the average number of years enrolled in school. In Brazil, maternal education was a categorical variable representing the percentage of mothers that had not completed primary school.

doi:10.1371/journal.pone.0047908.t002




Figure 1. Forest plot of the change in cognitive Z-score per 1% increase in diarrhea prevalence controlling for LAZ score as a continuous variable, mother's education, sex, age at cognition and SES.
doi:10.1371/journal.pone.0047908.g001

enabled us to control for residual confounding. In Philippines and Brazil, we observed a residual effect of age. In the Philippines, older children scored higher on the cognitive tests above and beyond what is accounted for in age-standardized test scores. However, in Brazil we observed a negative relationship between age at testing and cognitive performance. This may reflect increasing disparities over time such that children who demonstrate more advanced cognitive abilities early on and who live in stimulating environments continue to progress while children who have a lower skill level and are raised in non-stimulating environments fail to progress [22,23,24]. The gap between the two groups then widens as children age. In Brazil, diarrhea prevalence was found to be associated with lower cognitive test scores when the age variable was removed from the regression suggesting a correlation between age at testing and diarrhea prevalence (R. Pinkerton, personal communication). In Brazil this

might be due in part to progressive declines in early childhood diarrhea burdens during a decade of surveillance [25]. It appears that the effect of age on cognition is context dependent because it varies among these four studies. Socioeconomic conditions may confound the putative relationship between diarrhea and impaired cognition because poor or less educated families are likely to have both high rates of childhood diarrhea and delayed careseeking [26,27]. In impoverished households maternal and paternal education levels are often low and children receive less stimulation during times of health. It is likely that effects of diarrhea and/or stunting on cognition would be exacerbated in these deprived home environments during the diarrhea illness as stimulation would decline even further [28]. Studies have shown that the addition of stimulation provides an additional benefit to that of nutrition alone in improving mental development among stunted children [6]. Because cognitive function can be affected by both




Figure 2. Funnel plot analysis of possible bias in the 4 studies included in the meta-analysis of the beta coefficient for diarrhea prevalence.
doi:10.1371/journal.pone.0047908.g002

Table 3. Results Interpretation: Meta-analysis of the output from the regression of standardized cognitive z-score onto LAZ and diarrhea prevalence as continuous variables (model 1).^a.

Coefficient	Pooled Coefficient from Random Effects MA (95% CI)	p-value	Interpretation of Coefficient
% days with diarrhea	0.07 (−0.09, 0.23)	0.39	Cognitive z-score increased by 0.0695 per 1% increase in days with diarrhea among children, holding all other variables constant
LAZ continuous	0.18 (0.14, 0.21)	<0.001	Cognitive z-score increased by 0.1764 per Z-score increase in length for age in among children, holding all other variables constant
Sex	−0.10 (−0.18, −0.03)	0.004	Cognitive z-score was 0.1048 lower among males compared to females, holding all other variables constant

^a. We only report the pooled coefficients for variables with like definitions across study sites (i.e. diarrhea prevalence, LAZ, sex). Variables treated differently across study sites were not pooled (i.e. SES, age at cognitive assessment, maternal education).

doi:10.1371/journal.pone.0047908.t003

biological and environmental factors, studies that examine the possibility of confounding or interaction are the best to help us understand the possible role of diarrhea as a risk factor for cognitive impairment.

The complex pathways by which nutrition, infection, and the home environment affect a child's development are not well understood. Although there are limited data suggesting that numerous factors are at play, few studies have been able to look at all factors thus the roles of each remain elusive. Berkman et al observed that nutrition, parental education and school related factors independently affected cognition, but similar literature is scarce [19]. Our understanding of the complex relationships among diarrhea, nutritional status, poverty, careseeking, and developmental stimulation in the home would benefit greatly from additional studies explicitly designed to consider the role of each of these factors individually and more importantly to consider the mediating effects of one factor on another.

We did not demonstrate an association between diarrhea and cognition. Stunting remained a statistically significant covariate in this analysis showing a remarkably homogeneous association and suggesting a consistent relationship between stunting and cognition. While diarrhea may not directly impact cognitive impairment, the links between diarrhea and stunting, and stunting and

cognition are clear and provide additional rationale for accelerating interventions to reduce diarrhea.

Acknowledgments

We gratefully acknowledge the many key collaborators who made these studies possible, especially Dr. Aldo AM Lima and Dr. Sean Moore in Fortaleza, among many important field workers and collaborators.

Disclaimer: The views expressed in this article are those of the authors only and do not necessarily reflect the official policy or position of the Department of the Navy, Department of Defense, nor the U.S. Government.

Copyright statement: One author of this manuscript is an employee of the U.S. Government. This work was prepared as part of his duties. Title 17 U.S.C. § 105 provides that 'Copyright protection under this title is not available for any work of the United States Government.' Title 17 U.S.C. § 101 defines a U.S. Government work as a work prepared by a military service member or employee of the U.S. Government as part of that person's official duties.

Author Contributions

Conceived and designed the experiments: CLFW LA RLG AGL RM RP REB. Performed the experiments: RLG AGL RM RP LA. Analyzed the data: LL CFW. Wrote the paper: CLFW LL LA RLG AGL RM RP REB.

References

- Black RE, Cousens S, Johnson HL, Lawn JE, Rudan I, et al. (2010) Global, regional, and national causes of child mortality in 2008: a systematic analysis. *Lancet* 375: 1969–1987.
- Guerrant RL, Oria RB, Moore SR, Oria MO, Lima AA (2008) Malnutrition as an enteric infectious disease with long-term effects on child development. *Nutrition reviews* 66: 487–505.
- Guerrant DI, Moore SR, Lima AA, Patrick PD, Schorling JB, et al. (1999) Association of early childhood diarrhea and cryptosporidiosis with impaired physical fitness and cognitive function four-seven years later in a poor urban community in northeast Brazil. *The American journal of tropical medicine and hygiene* 61: 707–713.
- Niehaus MD, Moore SR, Patrick PD, Derr LL, Lorntz B, et al. (2002) Early childhood diarrhea is associated with diminished cognitive function 4 to 7 years later in children in a northeast Brazilian shantytown. *The American journal of tropical medicine and hygiene* 66: 590–593.
- Checkley W, Buckley G, Gilman RH, Assis AM, Guerrant RL, et al. (2008) Multi-country analysis of the effects of diarrhoea on childhood stunting. *International journal of epidemiology* 37: 816–830.
- Grantham-McGregor S (1995) A review of studies of the effect of severe malnutrition on mental development. *The Journal of nutrition* 125: 2233S–2238S.
- Mendez MA, Adair LS (1999) Severity and timing of stunting in the first two years of life affect performance on cognitive tests in late childhood. *The Journal of nutrition* 129: 1555–1562.
- Brown JL, Pollitt E (1996) Malnutrition, poverty and intellectual development. *Scientific American* 274: 38–43.
- Kaur A, Chowdhury S, Kumar R (1994) Mothers' beliefs and practices regarding prevention and management of diarrheal diseases. *Indian pediatrics* 31: 55–57.
- Guthrie GM, Butler A, Gorlow L, White GN (1964) Non-Verbal Expression of Self-Attitudes of Retardates. *American journal of mental deficiency* 69: 42–49.
- (1990) Manual for Test Nonverbal Intelligence. Austin: Pro-Ed.
- Wechsler D (1974) Intelligence scales for children-revised. San Antonio: The Psychological Corporation, Harcourt Brace Jovanovich.
- Pollitt E, Gorman KS, Engle PL, Martorell R, Rivera J (1993) Early supplementary feeding and cognition: effects over two decades. *Monographs of the Society for Research in Child Development* 58: 1–99; discussion 111–118.
- World Health Organization (2010) WHO child growth standards, SAS macros. Geneva: World Health Organization.
- Biostat (2005) Comprehensive Mets-Analysis Software Version 2. Englewood, NH: Biostat, Inc.
- StataCorp (2009) Stata Statistical Software Release 11. College Station, TX: StataCorp, L.P.
- Adair LS, Popkin BM, Akin JS, Guilkey DK, Gultiano S, et al. (2011) Cohort profile: the Cebu longitudinal health and nutrition survey. *International journal of epidemiology* 40: 619–625.
- Popkin BM, Adair L, Akin JS, Black R, Briscoe J, et al. (1990) Breast-feeding and diarrheal morbidity. *Pediatrics* 86: 874–882.
- Berkman DS, Lescano AG, Gilman RH, Lopez SL, Black MM (2002) Effects of stunting, diarrhoeal disease, and parasitic infection during infancy on cognition in late childhood: a follow-up study. *Lancet* 359: 564–571.
- Patrick PD, Oria RB, Madhavan V, Pinkerton RC, Lorntz B, et al. (2005) Limitations in verbal fluency following heavy burdens of early childhood

- diarrhea in Brazilian shantytown children. *Child neuropsychology : a journal on normal and abnormal development in childhood and adolescence* 11: 233–244.
21. Martorell R, Habicht JP, Rivera JA (1995) History and design of the INCAP longitudinal study (1969–77) and its follow-up (1983–89). *The Journal of nutrition* 125: 1027S–1041S.
 22. Noble KG, Tottenham N, Casey BJ (2005) Neuroscience perspectives on disparities in school readiness and cognitive achievement. *The Future of children/Center for the Future of Children, the David and Lucile Packard Foundation* 15: 71–89.
 23. Fernald LC, Weber A, Galasso E, Ratsifandrihamanana L (2011) Socioeconomic gradients and child development in a very low income population: evidence from Madagascar. *Developmental science* 14: 832–847.
 24. Burchinal M, McCartney K, Steinberg L, Crosnoe R, Friedman SL, et al. (2011) Examining the black-white achievement gap among low-income children using the NICHD study of early child care and youth development. *Child Development* 82: 1404–1420.
 25. Moore SR, Lima AA, Schorling JB, Barboza MS, Jr., Soares AM, et al. (2000) Changes over time in the epidemiology of diarrhea and malnutrition among children in an urban Brazilian shantytown, 1989 to 1996. *International journal of infectious diseases : IJID : official publication of the International Society for Infectious Diseases* 4: 179–186.
 26. Galler JR, Ramsey F, Solimano G, Lowell WE (1983) The influence of early malnutrition on subsequent behavioral development. II. Classroom behavior. *J Am Acad Child Psychiatry* 22: 16–22.
 27. Sigman M, McDonald MA, Neumann C, Bwibo N (1991) Prediction of cognitive competence in Kenyan children from Toddler nutrition, family characteristics and abilities. *J Child Psychol Psychiatry* 32: 307–320.
 28. Grantham-McGregor S (1995) A review of studies of the effect of severe malnutrition on mental development. *J Nutr* 125: 2233S–2238S.